

Matrices

CS5011 : Linear Algebra - basics

Matrix Arithmetic and Operation

- **Equality:** $A = B$ provided *dimensions of A and B are equal* and $a_{ij} = b_{ij}$ for all i and j .
Matrices of different sizes cannot be equal.
- **Addition, Subtraction:** $A_{n \times m} \pm B_{n \times m} = [a_{ij} \pm b_{ij}]$. *Matrices of different sizes cannot be added or subtracted.*
- **Scalar Multiple:** $cA = [ca_{ij}]$; c is any number.
- **Multiplication:** $A_{n \times p} * B_{p \times m} = A \cdot B_{n \times m}$
- **Transpose:** $A = [a_{ij}]_{n \times m}$ then $A^T = [a_{ji}]_{m \times n} \forall i, j$
- **Trace:** $tr(A) = \sum_{i=1}^n a_{ii}$. *If A is not square then trace is not defined.*

Properties of Matrix Arithmetic and the Transpose

- $A + B = B + A$
- $A + (B + C) = (A + B) + C$
- $A(BC) = (AB)C$
- $A(B \pm C) = AB \pm AC$
- $(B \pm C)A = BA \pm CA$
- $a(B \pm C) = aB \pm aC$
- $(a \pm b)C = aC \pm bC$
- $(ab)C = a(bC)$
- $a(BC) = (aB)C = B(aC)$
- $A(B) \neq B(A)$, in general.

Letters in caps define matrices, while that in small denote scalars.

Properties of Matrix Arithmetic and the Transpose

- $A + 0 = 0 + A = A$
- $A - A = 0$
- $0 - A = A$
- $0A = 0$ and $A0 = 0$
- $A^n A^m = A^{n+m}$
- $(A^n)^m = A^{nm}$
- $(A^T)^T = A$
- $(A \pm B)^T = A^T \pm B^T$
- $(cA)^T = cA^T$
- $(AB)^T = B^T A^T$

Letters in caps define matrices, while that in small denote scalars.

Important properties of the inverse matrix

Suppose that A and B are invertible matrices of the same size. Then,

a) AB is invertible and $(AB)^{-1} = B^{-1}A^{-1}$

b) A^{-1} is invertible and $(A^{-1})^{-1} = A$

c) For $n = 0, 1, 2, \dots$ A^n is invertible and $(A^n)^{-1} = A^{-n} = (A^{-1})^n$

d) If c is any non zero scalar then cA is invertible and $(cA)^{-1} = \frac{1}{c}A^{-1}$.

e) A^T is invertible and $(A^T)^{-1} = (A^{-1})^T$

Inverse Calculation

- The matrix $A = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$

will be ***invertible*** if $ad - bc \neq 0$

and ***singular*** if $ad - bc = 0$.

- If the matrix is invertible its inverse will be,

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}$$

Special Matrices : Diagonal Matrix

- **Diagonal Matrix:** A square matrix is called **diagonal** if it has the following form

$$D = \begin{bmatrix} d_1 & 0 & \dots & 0 \\ 0 & d_2 & \dots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & d_n \end{bmatrix}$$

- Suppose D is a diagonal matrix and $d_i, i = 1, \dots, n$ are the entries on the main diagonal.
- If one or more of the d_i 's are zero then the matrix is singular.

Diagonal Matrix (contd.)

- On the other hand if $d_i \neq 0, \forall i$ then the matrix is invertible and the inverse is,

$$D^{-1} = \begin{bmatrix} \frac{1}{d_1} & 0 & 0 & \dots & 0 \\ 0 & \frac{1}{d_2} & 0 & \dots & 0 \\ 0 & 0 & \frac{1}{d_3} & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & 0 & \frac{1}{d_n} \end{bmatrix}$$

Triangular matrix

$$U = \begin{bmatrix} u_{11} & u_{12} & u_{13} & \cdots & u_{1n} \\ 0 & u_{22} & u_{23} & \cdots & u_{2n} \\ 0 & 0 & u_{33} & \cdots & u_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & u_{nn} \end{bmatrix}_{n \times n}$$

Upper Triangular Matrix

$$L = \begin{bmatrix} l_{11} & 0 & 0 & \cdots & 0 \\ l_{21} & l_{22} & 0 & \cdots & 0 \\ l_{31} & l_{32} & l_{33} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ l_{n1} & l_{n2} & l_{n3} & \cdots & l_{nn} \end{bmatrix}_{n \times n}$$

Lower Triangular Matrix

- *If A is a triangular matrix with main diagonal entries $a_{11}, a_{22}, \dots, a_{nn}$ then if one or more of the a_{ii} 's are zero the matrix will be **singular**.*
- *On the other hand if $a_{ii} \neq 0 \forall i$ then the matrix is **invertible**.*

Symmetric and anti-symmetric matrices

Suppose that A is an $n \times m$ matrix, then A will be called **symmetric** if $A = A^T$.

Some properties of symmetric matrices are:

- a) *For any matrix A , both AA^T and $A^T A$ are symmetric.*
- b) *If A is an invertible symmetric matrix then A^{-1} is also symmetric.*
- c) *If A is invertible then AA^T and $A^T A$ are both invertible.*

Anti-Symmetric or Skew-Symmetric:

An anti-symmetric matrix is a square matrix that satisfies the identity $\mathbf{A} = -\mathbf{A}^T$.

Other Special forms of matrices:

- **Toeplitz matrix**
- **Block Circulant Matrix**
- **Orthogonal (also, -skew -sym)**
- **PD, PSD, ...**
- **Tri-diagonal system**
- **Hessian**
- **Jacobian**
- **Adjoint and Adjugate matrices**
- **(skew-) Hermitian (or self-adjoint) matrix**
- **Covariance matrix**
- **Periodic matrices**
- **Compound Matrix**
- **g-inv & Pseudo-inv**
- **GRAM matrix**
- **Kernel of matrix**
- **Schur Complement**
- **PERM (n)**
- **Skew-symmetric**
- **DFT Matrix**
- **Idempotent Matrices**
- **Vandermonde Matrices**

Matrix Multiplication

1. The order makes a difference...AB is different from BA.
2. **Rule** : The number of columns in first matrix must equal number of rows in second matrix.

In other words, the **inner dimensions** must be equal.

3. **Dimension of product** : The answer will be number of rows in first matrix by number of columns in second matrix.

In other words, the **outer dimensions**.

$$\begin{array}{c} \begin{bmatrix} 4 \\ 2 \end{bmatrix} \times \begin{bmatrix} 3 & 1 \end{bmatrix} = \begin{bmatrix} \square & \square \\ \square & \square \end{bmatrix} \\ \underbrace{2 \times 1 \quad 1 \times 2} \end{array}$$


$$\begin{array}{c} \begin{bmatrix} 3 & 1 \end{bmatrix} \times \begin{bmatrix} 4 \\ 2 \end{bmatrix} = \begin{bmatrix} \square \end{bmatrix} \\ \underbrace{1 \times 2 \quad 2 \times 1} \end{array}$$

Matrix Multiplication

Multiplication of two matrices, **A** and **B**, produces the matrix **C** whose elements, $c_{i,j}$ ($0 \leq i < n, 0 \leq j < m$), are computed as follows:

$$c_{i,j} = \sum_{k=0}^{l-1} a_{i,k} b_{k,j}$$

where **A** is an $n \times p$ matrix and **B** is an $p \times m$ matrix.



Matrix Notation and Matrix Multiplication

Nine co-efficients	$2u + v + w = 5$
Three unknowns	$4u - 6v = -2$
Three right-hand sides	$-2u + 7v + 2w = 9$

$$Ax = b$$

$$A = \begin{bmatrix} 2 & 1 & 1 \\ 4 & -6 & 0 \\ -2 & 7 & 2 \end{bmatrix}$$

Co-efficient matrix

$$x = \begin{bmatrix} u \\ v \\ w \end{bmatrix}$$

Solution vector

$$b = \begin{bmatrix} 5 \\ -2 \\ 9 \end{bmatrix}$$

constant vector

There are two ways to multiply a matrix A and a vector x .

- One way is a row at a time, each row of A combines with x to give a component of Ax . There are three inner products when A has three rows:

$$Ax \text{ by rows} \quad \begin{bmatrix} 1 & 1 & 6 \\ 3 & 0 & 1 \\ 1 & 1 & 4 \end{bmatrix} \begin{bmatrix} 2 \\ 5 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \cdot 2 + 1 \cdot 5 + 6 \cdot 0 \\ 3 \cdot 2 + 0 \cdot 5 + 3 \cdot 0 \\ 1 \cdot 2 + 1 \cdot 5 + 4 \cdot 0 \end{bmatrix} = \begin{bmatrix} 7 \\ 6 \\ 7 \end{bmatrix}$$

- Second way, multiplication a column at a time. The product Ax is found all at once, as a combination of the three columns of A :

$$Ax \text{ by columns} \quad 2 \begin{bmatrix} 1 \\ 3 \\ 1 \end{bmatrix} + 5 \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix} + 0 \begin{bmatrix} 6 \\ 3 \\ 4 \end{bmatrix} = \begin{bmatrix} 7 \\ 6 \\ 7 \end{bmatrix}$$

Properties of matrix multiplication


- Every product Ax can be found using whole columns. Therefore Ax is a combination of the columns of A . The coefficients are the components of x .
- The identity matrix I , with 1s on the diagonal and 0s everywhere else, leaves every vector unchanged.

Identity matrix $IA = A$ and $BI = B$.

Properties of matrix multiplication

- The i, j entry of AB is the inner product of the i -th row of A and the j -th column of B

$$(AB)_{32} = a_{31}b_{12} + a_{32}b_{22} + a_{33}b_{32} + a_{34}b_{42}$$



Properties of matrix multiplication

- Each entry of AB is the product of a row and a column:

$$(AB)_{ij} = (\text{row } i \text{ of } A) \text{ times } (\text{column } j \text{ of } B)$$

- Each column of AB is the product of a matrix and a column:

$$\text{column } j \text{ of } AB = A \text{ times } (\text{column } j \text{ of } B)$$

- Each row of AB is the product of a row and a matrix:

$$\text{row } i \text{ of } AB = (\text{row } i \text{ of } A) \text{ times } B$$

Properties of matrix multiplication

- For matrices A, B, C, D, E and F ,
- Matrix multiplication is **associative**:

$$(AB)C = A(BC)$$

- Matrix operations are **distributive**:

$$A(B + C) = AB + AC \text{ and } (B + C)D = BD + CD$$

- Matrix multiplication is **not commutative**: Usually
 $FE \neq EF$

Exception :

$$E = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \quad \text{and} \quad F = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} \quad EF = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 1 & 0 & 1 \end{bmatrix} = FE$$

Norms

To meter the lengths of vectors in a vector space we need the idea of a **norm**.

Norm is a function that maps x to a nonnegative real number

$$\| \cdot \|: F \rightarrow R^+$$

A Norm must satisfy following properties:

1 – Positivity $\|x\| > 0, \forall x \neq 0$

2 – Homogeneity $\|\alpha x\| = |\alpha| \|x\|, \forall x \in F \text{ and } \forall \alpha \in C$

3 – Triangle inequality $\|x + y\| \leq \|x\| + \|y\|, \forall x, y \in F$

Norm of vectors

p-norm is: $\|x\|_p = \left(\sum_i |a_i|^p \right)^{\frac{1}{p}} \quad p \geq 1$

For $p=1$ we have **1-norm** or **sum norm** $\|x\|_1 = \left(\sum_i |a_i| \right)$

For $p=2$ we have **2-norm** or **euclidian norm** $\|x\|_2 = \left(\sum_i |a_i|^2 \right)^{1/2}$

For $p=\infty$ we have **∞ -norm** or **max norm** $\|x\|_\infty = \max_i \{ |a_i| \}$

The l_p -Norm

The l_p - Norm for a vector x is defined as ($p \geq 1$):

$$\|x\|_{l_p} = \left(\sum_{i=1}^n |x_i|^p \right)^{1/p}$$

Examples:

- for $p=2$ we have the ordinary euclidian norm: $\|x\|_{l_2} = \sqrt{x^T x}$

- for $p= \infty$ the definition is $\|x\|_{l_\infty} = \max_{1 \leq i \leq n} |x_i|$

- a norm for matrices is induced via $\|A\| = \max_{x \neq 0} \frac{\|Ax\|}{\|x\|}$

- for l_2 this means :
 $\|A\|_2 = \text{maximum eigenvalue of } A^T A$

Properties of Matrix Norms

- These induced matrix norms satisfy:

$$\|A\| > 0 \text{ if } A \neq 0$$

$$\|\gamma A\| = |\gamma| \cdot \|A\| \text{ for any scalar } \gamma$$

$$\|A + B\| \leq \|A\| + \|B\| \text{ (triangle inequality)}$$

$$\|AB\| \leq \|A\| \cdot \|B\|$$

$$\|Ax\| \leq \|A\| \cdot \|x\| \text{ for any vector } x$$

Condition Number

- If A is square and nonsingular, then

$$\text{cond}(A) = \|A\| \cdot \|A^{-1}\|$$

- If A is singular, then $\text{cond}(A) = \infty$
- If A is nearly singular, then $\text{cond}(A)$ is large.
- The condition number measures the ratio of maximum stretch to maximum shrinkage:

$$\|A\| \cdot \|A^{-1}\| = \left(\max_{x \neq 0} \frac{\|Ax\|}{\|x\|} \right) \cdot \left(\min_{x \neq 0} \frac{\|Ax\|}{\|x\|} \right)^{-1}$$

The Gaussian Elimination Method

- The Gaussian elimination method is a **technique** for **solving systems of linear equations** of any size.
- The operations of the Gaussian elimination method are:
 1. **Interchange** any two equations.
 2. **Replace** an equation by a **nonzero constant multiple** of itself.
 3. **Replace** an equation by the **sum** of that equation and a **constant multiple of any other equation**.

Row-Reduced Form of a Matrix

- Each row consisting entirely of **zeros** lies **below** all rows having **nonzero entries**.
- The **first nonzero entry** in each nonzero row is **1** (called a **leading 1**).
- In any two successive (nonzero) rows, the **leading 1** in the lower row lies **to the right** of the **leading 1** in the **upper row**.
- If a column contains a **leading 1**, then the other entries in that column are **zeros**.

Row Operations

1. Interchange any two rows.
2. Replace any row by a nonzero constant multiple of itself.
3. Replace any row by the sum of that row and a constant multiple of any other row.

Terminology for the Gaussian Elimination Method

Unit Column

- A column in a coefficient matrix is in unit form if **one** of the entries in the column is a **1** and the **other** entries are **zeros**.

Pivoting

- The **sequence of row operations** that **transforms** a **given column** in an augmented matrix into a **unit column**.

Notation for Row Operations

- Letting R_i denote the i -th row of a matrix, we write

Operation 1: $R_i \leftrightarrow R_j$ to mean:
Interchange row i with row j .

Operation 2: cR_i to mean:
replace row i with c times row i .

Operation 3: $R_i + aR_j$ to mean:
Replace row i with the sum of row i and a times row j .

Example

- Pivot the matrix about the circled element

$$\begin{bmatrix} 3 & 5 & 9 \\ 2 & 3 & 5 \end{bmatrix}$$

Solution:

$$\begin{bmatrix} 3 & 5 & | & 9 \\ 2 & 3 & | & 5 \end{bmatrix} \xrightarrow{\frac{1}{3}R_1} \begin{bmatrix} 1 & 5/3 & | & 3 \\ 2 & 3 & | & 5 \end{bmatrix} \xrightarrow{R_2 - 2R_1} \begin{bmatrix} 1 & 5/3 & | & 3 \\ 0 & -1/3 & | & -1 \end{bmatrix}$$

The Gaussian Elimination Method

1. Write the **augmented matrix** corresponding to the linear system.
2. **Interchange rows**, if necessary, to obtain an augmented matrix in which the **first entry** in the **first row** is **nonzero**. Then **pivot** the matrix about this entry.
3. **Interchange** the **second row** with any row below it, if necessary, to obtain an augmented matrix in which the **second entry** in the **second row** is **nonzero**. **Pivot** the matrix about this entry.
4. **Continue** until the final matrix is in **row-reduced form**.

Augmented Matrices

- Matrices are **rectangular arrays of numbers** that can aid us by **eliminating the need to write the variables** at each step of the reduction.
- For example, the **system**

$$2x + 4y + 6z = 22$$

$$3x + 8y + 5z = 27$$

$$-x + y + 2z = 2$$

may be represented by the **augmented matrix**

$$\left[\begin{array}{ccc|c} 2 & 4 & 6 & 22 \\ 3 & 8 & 5 & 27 \\ -1 & 1 & 2 & 2 \end{array} \right]$$

Augmented matrix
[C|B]

Coefficient
Matrix [C]

Matrices and Gaussian Elimination

- **Every step** in the **Gaussian elimination method** can be expressed with **matrices**, rather than systems of equations, thus simplifying the whole process:

- Steps expressed as **systems of equations**:

$$2x + 4y + 6z = 22$$

$$3x + 8y + 5z = 27$$

$$-x + y + 2z = 2$$

- Steps expressed as **augmented matrices**:

$$\left[\begin{array}{ccc|c} 2 & 4 & 6 & 22 \\ 3 & 8 & 5 & 27 \\ -1 & 1 & 2 & 2 \end{array} \right]$$

$$\begin{aligned} 2x + 4y + 6z &= 22 \\ 3x + 8y + 5z &= 27 \\ -x + y + 2z &= 2 \end{aligned}$$

$$\begin{aligned} x + 2y + 3z &= 11 \\ 3x + 8y + 5z &= 27 \\ -x + y + 2z &= 2 \end{aligned}$$

$$\begin{aligned} x + 2y + 3z &= 11 \\ 2y - 4z &= -6 \\ -x + y + 2z &= 2 \end{aligned}$$


$$\begin{aligned} x + 2y + 3z &= 11 \\ 2y - 4z &= -6 \\ 3y + 5z &= 13 \end{aligned}$$

$$\left[\begin{array}{ccc|c} 2 & 4 & 6 & 22 \\ 3 & 8 & 5 & 27 \\ -1 & 1 & 2 & 2 \end{array} \right]$$


$$\left[\begin{array}{ccc|c} 1 & 2 & 3 & 11 \\ 3 & 8 & 5 & 27 \\ -1 & 1 & 2 & 2 \end{array} \right]$$

$$\left[\begin{array}{ccc|c} 1 & 2 & 3 & 11 \\ 0 & 2 & -4 & -6 \\ -1 & 1 & 2 & 2 \end{array} \right]$$


$$\left[\begin{array}{ccc|c} 1 & 2 & 3 & 11 \\ 0 & 2 & -4 & -6 \\ 0 & 3 & 5 & 13 \end{array} \right]$$




$$R'_1 = \frac{1}{2}R_1$$



$$R'_2 = R_2 - 3R_1$$



$$R'_3 = R_3 + R_1$$



$$R'_2 = \frac{1}{2}R_2$$

$$\begin{aligned}x + 2y + 3z &= 11 \\y - 2z &= -3 \\3y + 5z &= 13\end{aligned}$$

$$\begin{aligned}x + 7z &= 11 \\y - 2z &= -3 \\3y + 5z &= 13\end{aligned}$$

$$\begin{aligned}x + 7z &= 11 \\y - 2z &= -3 \\11z &= 22\end{aligned}$$


$$\begin{aligned}x + 7z &= 11 \\y - 2z &= -3 \\z &= 2\end{aligned}$$

$$\begin{bmatrix} 1 & 2 & 3 & 11 \\ 0 & 1 & -2 & -3 \\ 0 & 3 & 5 & 13 \end{bmatrix}$$


$$\begin{bmatrix} 1 & 0 & 7 & 17 \\ 0 & 1 & -2 & -3 \\ 0 & 3 & 5 & 13 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 7 & 17 \\ 0 & 1 & -2 & -3 \\ 0 & 0 & 11 & 22 \end{bmatrix}$$


$$\begin{bmatrix} 1 & 0 & 7 & 17 \\ 0 & 1 & -2 & -3 \\ 0 & 0 & 1 & 2 \end{bmatrix}$$




$$R'_2 = \frac{1}{2}R_2$$




$$R'_1 = R_1 - 2R_2$$



$$R'_3 = R_3 - 3R_2$$




$$R'_3 = \frac{1}{11}R_3$$




$$R'_1 = R_1 - 7R_3$$

$$\begin{array}{l}
 x = 3 \\
 y - 2z = -3 \\
 z = 2
 \end{array}
 \quad \left| \begin{array}{ccc|c}
 1 & 0 & 0 & 3 \\
 0 & 1 & -2 & -3 \\
 0 & 0 & 1 & 2
 \end{array} \right.$$

 $R'_1 = R_1 - 7R_3$

$$\begin{array}{l}
 x = 3 \\
 y = -3 \\
 z = 2
 \end{array}
 \quad \left| \begin{array}{ccc|c}
 1 & 0 & 0 & 3 \\
 0 & 1 & 0 & 1 \\
 0 & 0 & 1 & 2
 \end{array} \right.$$

 $R'_2 = R_2 + 2R_3$

**Row Reduced Form
of the Matrix**

Thus, the **solution** to the system is $x = 3$, $y = 1$,
and $z = 2$.

Gaussian Elimination in the case of unique solution

- With a full set of n pivots, there is only one solution.
- The system is non singular, and it is solved by forward elimination and back-substitution.

Systems with no Solution

- If there is a **row** in the augmented matrix containing **all zeros** to the **left** of the **vertical line** and a **nonzero** entry to the **right** of the **line**, then the system of equations has **no solution**.

Theorem

- a. If the **number of equations** is **greater (over-determined system)** than or equal to the **number of variables** in a linear system, then one of the following is true:
 - i. The system has **no solution**.
 - ii. The system has **exactly one solution**.
 - iii. The system has **infinitely many solutions**.

- b. If there are **fewer equations than variables (under-determined system)** in a linear system, then the system either has **no solution** or it has **infinitely many solutions**.

Inverse matrix

- The inverse of an n by n matrix is another n by n matrix. The inverse of A is written A^{-1} (and pronounced “ A inverse”).
- The fundamental property is simple: If you multiply by A and then multiply by A^{-1} , you are back where you started:

Inverse matrix If $b = Ax$ then $A^{-1}b = x$

- Thus $A^{-1}Ax = x$. The matrix A^{-1} times A is the identity matrix. ***Not all matrices have inverses. An inverse is impossible when Ax is zero and x is nonzero.*** Then A^{-1} would have to get back from $Ax = 0$ to x . No matrix can multiply that zero vector Ax and produce a nonzero vector x .
- Our goals are to define the inverse matrix and compute it and use it, when A^{-1} exists—and then to understand which matrices don’t have inverses.

Properties : Inverse matrix

1K The **inverse** of A is a matrix B such that $BA = I$ and $AB = I$. There is at most one such B , and it is denoted by A^{-1} :

$$A^{-1}A = I \quad \text{and} \quad AA^{-1} = I. \quad (1)$$

Note 1. *The inverse exists if and only if elimination produces n pivots* (row exchanges allowed). Elimination solves $Ax = b$ without explicitly finding A^{-1} .

Note 2. The matrix A cannot have two different inverses, Suppose $BA = I$ and also $AC = I$. Then $B = C$, according to this “proof by parentheses”:

$$B(AC) = (BA)C \quad \text{gives} \quad BI = IC \quad \text{which is} \quad B = C. \quad (2)$$

This shows that a *left-inverse* B (multiplying from the left) and a *right-inverse* C (multiplying A from the right to give $AC = I$) must be the *same matrix*.

Note 3. If A is invertible, the one and only solution to $Ax = b$ is $x = A^{-1}b$:

$$\textbf{Multiply} \quad Ax = b \quad \textbf{by} \quad A^{-1}. \quad \textbf{Then} \quad x = A^{-1}Ax = A^{-1}b.$$

Note 4. (Important) *Suppose there is a nonzero vector x such that $Ax = 0$. Then A cannot have an inverse.* To repeat: No matrix can bring 0 back to x .

If A is invertible, then $Ax = 0$ can only have the zero solution $x = 0$.

Properties : Inverse matrix

Note 5. A 2 by 2 matrix is invertible if and only if $ad - bc$ is not zero:

$$\text{2 by 2 inverse} \quad \begin{bmatrix} a & b \\ c & d \end{bmatrix}^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}. \quad (3)$$

This number $ad - bc$ is the *determinant* of A . A matrix is invertible if its determinant is not zero (Chapter 4). In **MATLAB**, the invertibility test is *to find n nonzero pivots*. Elimination produces those pivots before the determinant appears.

Note 6. A diagonal matrix has an inverse provided no diagonal entries are zero:

$$\text{If } A = \begin{bmatrix} d_1 & & \\ & \ddots & \\ & & d_n \end{bmatrix} \text{ then } A^{-1} = \begin{bmatrix} 1/d_1 & & \\ & \ddots & \\ & & 1/d_n \end{bmatrix} \text{ and } AA^{-1} = I.$$

When two matrices are involved, not much can be done about the inverse of $A + B$. The sum might or might not be invertible. Instead, it is the inverse of their *product* AB which is the key formula in matrix computations. Ordinary numbers are the same: $(a + b)^{-1}$ is hard to simplify, while $1/ab$ splits into $1/a$ times $1/b$. But for matrices *the order of multiplication must be correct*—if $ABx = y$ then $Bx = A^{-1}y$ and $x = B^{-1}A^{-1}y$. **The inverses come in reverse order.**

Properties : Inverse matrix

1L A product AB of invertible matrices is inverted by $B^{-1}A^{-1}$:

$$\text{Inverse of } AB \quad (AB)^{-1} = B^{-1}A^{-1}. \quad (4)$$

Proof. To show that $B^{-1}A^{-1}$ is the inverse of AB , we multiply them and use the associative law to remove parentheses. Notice how B sits next to B^{-1} :

$$(AB)(B^{-1}A^{-1}) = ABB^{-1}A^{-1} = AIA^{-1} = AA^{-1} = I$$

$$(B^{-1}A^{-1})(AB) = B^{-1}A^{-1}AB = B^{-1}IB = B^{-1}B = I.$$

□

A similar rule holds with three or more matrices:

$$\text{Inverse of } ABC \quad (ABC)^{-1} = C^{-1}B^{-1}A^{-1}.$$

We saw this change of order when the elimination matrices E , F , G were inverted to come back from U to A . In the forward direction, $GFEA$ was U . In the backward direction, $L = E^{-1}F^{-1}G^{-1}$ was the product of the inverses. *Since G came last, G^{-1} comes first.* Please check that A^{-1} would be $U^{-1}GFE$.

Calculation of A^{-1} : The Gauss-Jordan Method

- Given the $n \times n$ matrix A :
 1. Adjoin the $n \times n$ identity matrix I to obtain the augmented matrix $[A | I]$.
 2. Use a sequence of row operations to reduce $[A | I]$ to the form $[I | B]$ if possible.
- Then the matrix B is the inverse of A .

Example

- Find the inverse of the matrix $A = \begin{bmatrix} 2 & 1 & 1 \\ 3 & 2 & 1 \\ 2 & 1 & 2 \end{bmatrix}$

Solution

- We form the **augmented matrix**

$$\left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 3 & 2 & 1 & 0 & 1 & 0 \\ 2 & 1 & 2 & 0 & 0 & 1 \end{array} \right]$$

Example

- Find the inverse of the matrix $A = \begin{bmatrix} 2 & 1 & 1 \\ 3 & 2 & 1 \\ 2 & 1 & 2 \end{bmatrix}$

Solution

- And use the **Gauss-Jordan elimination method** to **reduce it** to the form $[I | B]$:

$$\left[\begin{array}{ccc|ccc} 2 & 1 & 1 & 1 & 0 & 0 \\ 3 & 2 & 1 & 0 & 1 & 0 \\ 2 & 1 & 2 & 0 & 0 & 1 \end{array} \right] \xrightarrow{R_1 - R_2} \left[\begin{array}{ccc|ccc} -1 & -1 & 0 & 1 & -1 & 0 \\ 3 & 2 & 1 & 0 & 1 & 0 \\ 2 & 1 & 2 & 0 & 0 & 1 \end{array} \right]$$

$$\left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 2 & -1 & 0 \\ 0 & 1 & -1 & -3 & 2 & 0 \\ 0 & 0 & 1 & -1 & 0 & 1 \end{array} \right] \xrightarrow{\begin{array}{l} R_1 + R_2 \\ -R_2 \\ R_3 - R_2 \end{array}} \left[\begin{array}{ccc|ccc} 1 & 1 & 0 & -1 & 1 & 0 \\ 0 & -1 & 1 & 3 & -2 & 0 \\ 0 & -1 & 2 & 2 & -2 & 1 \end{array} \right] \xrightarrow{\begin{array}{l} -R_1 \\ R_2 + 3R_3 \\ R_3 + 2R_1 \end{array}}$$

Example

- Find the inverse of the matrix $A = \begin{bmatrix} 2 & 1 & 1 \\ 3 & 2 & 1 \\ 2 & 1 & 2 \end{bmatrix}$

Solution

- And use the **Gauss-Jordan elimination method** to reduce it to the form $[I | B]$:

$$\begin{array}{c} \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 2 & -1 & 0 \\ 0 & 1 & -1 & -3 & 2 & 0 \\ 0 & 0 & 1 & -1 & 0 & 1 \end{array} \right] \xrightarrow{\substack{R_1 - R_3 \\ R_2 + R_3}} \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 3 & -1 & -1 \\ 0 & 1 & 0 & -4 & 2 & 1 \\ 0 & 0 & 1 & -1 & 0 & 1 \end{array} \right] \\ \text{Previous step} \qquad \qquad \underbrace{\hspace{10em}}_{I_n} \qquad \underbrace{\hspace{10em}}_B \end{array}$$

$$B = A^{-1} = \begin{bmatrix} 3 & -1 & -1 \\ -4 & 2 & 1 \\ -1 & 0 & 1 \end{bmatrix}$$

Finding the inverse of a square matrix using LU decomposition

The inverse $[B]$ of a square matrix $[A]$ is defined as

How can LU Decomposition be used to find the inverse?

Assume the first column of $[B]$ to be $[b_{11} \ b_{12} \ \dots \ b_{n1}]^T$
Using this and the definition of matrix multiplication

Example: Inverse of a Matrix

Find the inverse of a square matrix $[A]$

$$[A] = \begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix}$$

Using the decomposition procedure, the $[L]$ and $[U]$ matrices are found to be

$$[A] = [L][U] = \begin{bmatrix} 1 & 0 & 0 \\ 2.56 & 1 & 0 \\ 5.76 & 3.5 & 1 \end{bmatrix} \begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & 0 & 0.7 \end{bmatrix}$$

Example: Inverse of a Matrix

Solving for the each column of $[B]$ requires two steps

1) Solve $[L][Z] = [C]$ for $[Z]$

2) Solve $[U][X] = [Z]$ for $[X]$

$$\text{Step 1: } [L][Z] = [C] \rightarrow \begin{bmatrix} 1 & 0 & 0 \\ 2.56 & 1 & 0 \\ 5.76 & 3.5 & 1 \end{bmatrix} \begin{bmatrix} z_1 \\ z_2 \\ z_3 \end{bmatrix} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$$

This generates the equations:

$$z_1 = 1$$
$$2.56z_1 + z_2 = 0$$
$$5.76z_1 + 3.5z_2 + z_3 = 0$$

Example: Inverse of a Matrix

Solving for $[Z]$

$$z_1 = 1$$

$$z_2 = 0 - 2.56z_1$$

$$= 0 - 2.56(1)$$

$$= -2.56$$

$$z_3 = 0 - 5.76z_1 - 3.5z_2$$

$$= 0 - 5.76(1) - 3.5(-2.56)$$

$$= 3.2$$

$$[Z] = \begin{bmatrix} z_1 \\ z_2 \\ z_3 \end{bmatrix} = \text{[Blue Box]}$$

Example: Inverse of a Matrix

Solving $[U][X] = [Z]$ for $[X]$

$$\begin{bmatrix} 25 & 5 & 1 \\ 0 & -4.8 & -1.56 \\ 0 & 0 & 0.7 \end{bmatrix} \begin{bmatrix} b_{11} \\ b_{21} \\ b_{31} \end{bmatrix} = \begin{bmatrix} 1 \\ -2.56 \\ 3.2 \end{bmatrix}$$

$$25b_{11} + 5b_{21} + b_{31} = 1$$

$$-4.8b_{21} - 1.56b_{31} = -2.56$$

$$0.7b_{31} = 3.2$$

Example: Inverse of a Matrix

Using Backward Substitution


$$b_{31} = \frac{3.2}{0.7} = 4.571$$

$$\begin{aligned} b_{21} &= \frac{-2.56 + 1.560b_{31}}{-4.8} \\ &= \frac{-2.56 + 1.560(4.571)}{-4.8} = -0.9524 \end{aligned}$$

$$\begin{aligned} b_{11} &= \frac{1 - 5b_{21} - b_{31}}{25} \\ &= \frac{1 - 5(-0.9524) - 4.571}{25} = 0.04762 \end{aligned}$$

So the first column of the inverse of

$$\begin{bmatrix} b_{11} \\ b_{21} \\ b_{31} \end{bmatrix} =$$



Example: Inverse of a Matrix

Repeating for the second and third columns of the inverse

Second Column

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix} \begin{bmatrix} b_{12} \\ b_{22} \\ b_{32} \end{bmatrix} = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix}$$

$$\begin{bmatrix} b_{12} \\ b_{22} \\ b_{32} \end{bmatrix} = \begin{bmatrix} \\ \\ \end{bmatrix}$$

Third Column

$$\begin{bmatrix} 25 & 5 & 1 \\ 64 & 8 & 1 \\ 144 & 12 & 1 \end{bmatrix} \begin{bmatrix} b_{13} \\ b_{23} \\ b_{33} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

$$\begin{bmatrix} b_{13} \\ b_{23} \\ b_{33} \end{bmatrix} = \begin{bmatrix} \\ \\ \end{bmatrix}$$

Example: Inverse of a Matrix

The inverse of $[A]$ is

$$[A]^{-1} = \begin{bmatrix} 0.04762 & -0.08333 & 0.03571 \\ -0.9524 & 1.417 & -0.4643 \\ 4.571 & -5.000 & 1.429 \end{bmatrix}$$

To check your work do the following operation

$$[A][A]^{-1} = [I] = [A]^{-1}[A]$$

Eigenvalues and Eigenvectors

CS6015/LARP

Ack: Linear Algebra and Its Applications , Gilbert Strang

The Solution of $Ax = \lambda x$

- $Ax = \lambda x$ is a nonlinear equation; λ multiplies x . If we could discover λ , then the equation for x would be **linear**.
- We could write λIx in place of λx , and bring this term over to the left side:

$$(A - \lambda I)x = 0$$

The vector x is in the nullspace of $A - \lambda I$.

The number λ is chosen so that $A - \lambda I$ has a nullspace.

- We want a **nonzero** eigenvector x . The vector $x = 0$ always satisfies $Ax = \lambda x$, but it is useless.
- To be of any use, the nullspace of $A - \lambda I$ must contain vectors other than zero.
- In short, **$A - \lambda I$ must be singular.**

The Solution of $Ax = \lambda x$

5A The number λ is an eigenvalue of A if and only if $A - \lambda I$ is singular:

$$\det(A - \lambda I) = 0. \quad (10)$$

This is the characteristic equation. Each λ is associated with eigenvectors x :

$$(A - \lambda I)x = 0 \quad \text{or} \quad Ax = \lambda x. \quad (11)$$

The Solution of $Ax = \lambda x$

- Example:

$$A = \begin{bmatrix} 4 & -5 \\ 2 & -3 \end{bmatrix} \quad \text{we shift } A \text{ by } \lambda I \text{ to make it singular:}$$

Subtract λI $A - \lambda I = \begin{bmatrix} 4 - \lambda & -5 \\ 2 & -3 - \lambda \end{bmatrix}$

Determinant $|A - \lambda I| = (4 - \lambda)(-3 - \lambda) + 10$ or $\lambda^2 - \lambda - 2$

- This is the **characteristic polynomial**.
- Its **roots**, where the **determinant is zero**, are the **eigenvalues**.

$$\lambda^2 - \lambda - 2 = (\lambda + 1)(\lambda - 2)$$

The Solution of $Ax = \lambda x$

Eigenvalues $\lambda = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{1 \pm \sqrt{9}}{2} = -1 \text{ and } 2.$

- There are two eigen values, because a quadratic has two roots.
- The values $\lambda = -1$ and $\lambda = 2$ lead to a solution of $Ax = \lambda x$ or $(A - \lambda I)x = 0$.

$$\lambda_1 = -1 : \quad (A - \lambda_1 I)x = \begin{bmatrix} 5 & -5 \\ 2 & -2 \end{bmatrix} \begin{bmatrix} y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The solution (the first eigenvector) is any nonzero multiple of x_1 :

Eigenvector for λ_1 $x_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$

The Solution of $Ax = \lambda x$

The solution (the first eigenvector) is any nonzero multiple of x_1 :

$$\text{Eigenvector for } \lambda_1 \quad x_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}.$$

The computation for λ_2 is done separately:

$$\lambda_2 = 2: \quad (A - \lambda_2 I)x = \begin{bmatrix} 2 & -5 \\ 2 & -5 \end{bmatrix} \begin{bmatrix} y \\ z \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

The second eigenvector is any nonzero multiple of x_2 :

$$\text{Eigenvector for } \lambda_2 \quad x_2 = \begin{bmatrix} 5 \\ 2 \end{bmatrix}.$$

The Solution of $Ax = \lambda x$

- The steps in solving $Ax = \lambda x$:
 - 1. Compute the determinant of $A - \lambda I$.** With λ subtracted along the diagonal, this determinant is a polynomial of degree n . It starts with $(-\lambda)^n$.
 - 2. Find the roots of this polynomial.** The n roots are the eigenvalues of A .
 - 3. For each eigenvalue solve the equation $(A - \lambda I)x = 0$.** Since the determinant is zero, there are solutions other than $x = 0$. Those are the eigenvectors.

The Solution of $Ax = \lambda x$ (Recap)

- The key equation was $Ax = \lambda x$.
- Most vectors x will not satisfy such an equation.
- They **change direction** when multiplied by A , so that Ax is not a multiple of x .
- ***This means that only certain special numbers are eigenvalues, and only certain special vectors x are eigenvectors.***

Example 3. The eigenvalues are on the main diagonal when A is triangular.

$$\det(A - \lambda I) = \begin{vmatrix} 1 - \lambda & 4 & 5 \\ 0 & \frac{3}{4} - \lambda & 6 \\ 0 & 0 & \frac{1}{2} - \lambda \end{vmatrix} = (1 - \lambda)\left(\frac{3}{4} - \lambda\right)\left(\frac{1}{2} - \lambda\right)$$

- The determinant is just the product of the diagonal entries.
- It is zero if $\lambda = 1, \lambda = \frac{3}{4},$ or $\lambda = \frac{1}{2}$
- The eigenvalues were already sitting along the main diagonal.

5B The *sum* of the n eigenvalues equals the sum of the n diagonal entries:

$$\text{Trace of } A = \lambda_1 + \cdots + \lambda_n = a_{11} + \cdots + a_{nn}. \quad (15)$$

Furthermore, the *product* of the n eigenvalues equals the *determinant* of A .

For a 2 by 2 matrix, the trace and determinant tell us everything:

$$\begin{bmatrix} a & b \\ c & d \end{bmatrix} \text{ has trace } a + d, \text{ and determinant } ad - bc$$

$$\det(A - \lambda I) = \det \begin{vmatrix} a - \lambda & b \\ c & d - \lambda \end{vmatrix} = \lambda^2 \text{ [redacted]}$$

$$\text{The eigenvalues are } \lambda = \text{[redacted]}^{1/2}.$$

Singular Value Decomposition

$A = U\Sigma V^T$ is known as the “**SVD**” or the ***singular value decomposition***.

The SVD is closely associated with the eigenvalue-eigenvector factorization $Q\Lambda Q^T$ of a positive definite matrix.

Any $m \times n$ matrix A can be factored into

$$A = U\Sigma V^T = (\mathbf{orthogonal})(\mathbf{diagonal})(\mathbf{orthogonal}).$$

The columns of U ($m \times m$) are ***eigenvectors of AA^T*** , and the columns of V ($n \times n$) are ***eigenvectors of $A^T A$*** .

The r singular values on the diagonal of Σ ($m \times n$) are the ***square roots of the nonzero eigenvalues*** of both AA^T and $A^T A$.

Singular Value Decomposition

Remark 1.

- For positive definite matrices, Σ is Λ and $U\Sigma V^T$ is identical to $Q\Lambda Q^T$.
- For other symmetric matrices, any negative eigenvalues in Λ become positive in Σ .
- For complex matrices, Σ remains real but U and V become *unitary* (the complex version of orthogonal).

Remark 2.

U and V give orthonormal bases for all four fundamental subspaces:

first	r	columns of U :	column space of A
last	$m - r$	columns of U :	left nullspace of A
first	r	columns of V :	row space of A
last	$n - r$	columns of V :	nullspace of A

Singular Value Decomposition

Remark 3.

Eigenvectors of AA^T and $A^T A$ must go into the columns of U and V :

$$AA^T = (U\Sigma V^T)(V\Sigma^T U^T) = U\Sigma\Sigma^T U^T \quad \text{and, similarly,} \quad A^T A = V\Sigma^T \Sigma V^T.$$

- U must be the eigenvector matrix for AA^T .
- The eigenvalue matrix in the middle is $\Sigma\Sigma^T$ — which is $m \times m$ with $\sigma_1^2, \dots, \sigma_r^2$ on the diagonal.
- From the $A^T A = V\Sigma^T \Sigma V^T$, the V matrix must be the eigenvector matrix for $A^T A$.

Singular Value Decomposition

Example 1.

This A has only one column: rank $r = 1$. Then Σ has only $\sigma_1 = 3$:

$$\mathbf{SVD} \quad A = \begin{bmatrix} -1 \\ 2 \\ 2 \end{bmatrix} = \begin{bmatrix} -\frac{1}{3} & \frac{2}{3} & \frac{2}{3} \\ \frac{2}{3} & -\frac{1}{3} & \frac{2}{3} \\ \frac{2}{3} & \frac{2}{3} & -\frac{1}{3} \end{bmatrix} \begin{bmatrix} 3 \\ 0 \\ 0 \end{bmatrix} \begin{bmatrix} 1 \end{bmatrix} = U_{3 \times 3} \Sigma_{3 \times 1} V_{1 \times 1}^T$$

$A^T A$ is 1 by 1, whereas AA^T is 3 by 3. They both have eigenvalue 9 (whose square root is the 3 in Σ). The two zero eigenvalues of AA^T leave some freedom for the eigenvectors in columns 2 and 3 of U . We kept that matrix orthogonal.

Singular Value Decomposition

Example 2.

Now A has rank 2, and $AA^T = \begin{bmatrix} 2 & -1 \\ -1 & 2 \end{bmatrix}$ with $\lambda = 3$ and 1:

$$\begin{bmatrix} -1 & 1 & 0 \\ 0 & -1 & 1 \end{bmatrix} = U\Sigma V^T = \frac{1}{\sqrt{2}} \begin{bmatrix} -1 & 1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} \sqrt{3} & 0 & 0 \\ 0 & 1 & 0 \end{bmatrix} \begin{bmatrix} 1 & -2 & 1 \\ -1 & 0 & 1 \\ 1 & 1 & 1 \end{bmatrix} \begin{matrix} /\sqrt{6} \\ / \sqrt{2} \\ / \sqrt{3} \end{matrix}$$

Notice $\sqrt{3}$ and $\sqrt{1}$. The columns of U are *left singular vectors* (unit eigenvectors of AA^T).

The columns of V are *right singular vectors* (unit eigenvectors of $A^T A$).

Applications of Singular Value Decomposition

Image Processing.

- Suppose a satellite takes a picture, and wants to send it to Earth.
- The picture may contain 1000×1000 “pixels” —a million little squares, each with a definite color.
- We can code the colors, and send back 1,000,000 numbers.
- It is ***better to find the essential information inside the 1000×1000 matrix***, and send only that.

In SVD some σ 's are significant and others are extremely small.

If we keep 20 and throw away 980, then we send only the corresponding 20 columns of U and V .

The other 980 columns are multiplied in $U\Sigma V^T$ by the small σ 's that are being ignored. ***If only 20 terms are kept, we send 20 times 2000 numbers instead of a million (25 to 1 compression).***

