

---

CS6015 : Linear Algebra and Random Processes  
Tutorial #2

Deadline: None

---

- This tutorial covers topics already covered in class. [Lecture slide 4-7]
  - While this is optional, it is strongly recommended that students solve this tutorial.
- 

NAME :

ROLL NUMBER :

1. Find if  $A = \begin{bmatrix} 1 & 2 & 3 & 4 \\ 0 & 2 & 3 & 4 \\ 0 & 0 & 3 & 4 \\ 0 & 0 & 0 & 4 \end{bmatrix}$  is invertible and compute  $A^{-1}$  by **Gauss-Jordan** method if it exists.

**Solution:**

2. For which right hand sides (find a condition on  $b_1, b_2, b_3$ ) are these systems solvable? Solve using **Gaussian Elimination**.

(a)  $\begin{bmatrix} 1 & 4 & 2 \\ 2 & 8 & 4 \\ -1 & -4 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$

(b)  $\begin{bmatrix} 1 & 4 \\ 2 & 9 \\ -1 & -4 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix}$

**Solution:**

3. Describe the column spaces (lines or planes) of the following two matrices.

$$A = \begin{bmatrix} 1 & 0 \\ 2 & 0 \\ 0 & 0 \end{bmatrix}, B = \begin{bmatrix} 1 & 0 \\ 0 & 2 \\ 0 & 0 \end{bmatrix}$$

**Solution:**

4. Comment on each of the following collection of vectors (in  $R^3$ ) and state if they are linearly independent.

(a)  $\left\{ \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} \right\}$

(d)  $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ 3 \\ 4 \end{bmatrix} \right\}$

(b)  $\left\{ \begin{bmatrix} 2 \\ 1 \\ 6 \end{bmatrix}, \begin{bmatrix} 5 \\ 2 \\ 2 \end{bmatrix}, \begin{bmatrix} 1 \\ 2 \\ 9 \end{bmatrix} \right\}$

(e)  $\left\{ \begin{bmatrix} 1 \\ 0 \\ 2 \end{bmatrix}, \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}, \begin{bmatrix} 17 \\ 0 \\ 0 \end{bmatrix} \right\}$

(c)  $\left\{ \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \right\}$

**Solution:**

5. Whether the following set of vectors span the entire  $R^3$ ? If not, then what does the span of these vectors represent?

$$\left\{ \begin{bmatrix} 1 \\ -1 \\ 4 \end{bmatrix}, \begin{bmatrix} -2 \\ 1 \\ 3 \end{bmatrix}, \begin{bmatrix} 4 \\ -3 \\ 5 \end{bmatrix} \right\}$$

**Solution:**

6. The span of vectors is the set of all their linear combinations. Consider two 2-D vectors  $v$  and  $w$ . What will be the spans of these vectors in the following cases:
- (a)  $v$  and  $w$  lie on the same line
  - (b)  $v$  and  $w$  do not lie on the same line
  - (c)  $v$  and  $w$  are zero

**Solution:**

7. Comment on the following set of transformations and state if they are linear transformations or not. The input is of the form,  $v = (v_1, v_2)$ .

(a)  $T(v) = (v_2, v_1)$

(c)  $T(v) = (0, v_1)$

(b)  $T(v) = (v_1, v_1)$

(d)  $T(v) = (0, 1)$

**Solution:**

8. Which of the following subsets of  $R^3$  are actually subspaces?

(a) The plane of vectors  $(b_1, b_2, b_3)$  with  $b_1 = b_2$

(b) The plane of vectors with  $b_1 = 1$

(c) The vectors with  $b_1 b_2 b_3 = 0$

(d) All linear combinations of  $v = [1, 4, 0]$  and  $w = [2, 2, 2]$

(e) All vectors that satisfy  $b_1 + b_2 + b_3 = 0$

(f) All vectors with  $b_1 \leq b_2 \leq b_3$

**Solution:**

9. How many  $(0, 1, \infty)$  solutions does this system of linear equation have? Write the specific answer if there exists a unique solution, or write the closed form expression if infinitely many solutions exist for this system of equations.

$$x_1 + x_3 + x_4 = 1$$

$$2x_2 + x_3 + x_4 = 0$$

$$x_1 + 2x_2 + x_3 = 1$$

**Solution:**

10. For a given set of vectors  $(v = [v_1, v_2, v_3 \dots v_n])$  in a vector space S, prove that the span(v) is a subspace of S.

**Solution:**

11. Prove that a square matrix can have at most one inverse.

**Solution:**

12. Mark the following statements as true/false:

- (a) The columns of a matrix are a basis for the column space.
- (b) In  $Ax = v$ , let  $x_1, x_2$  and  $v_1, v_2$  be the basis vectors of the input and output spaces respectively. In the input space, if a vector is  $= c(x_1) + d(x_2)$ , then the transformed vector in output space will be  $= c(v_1) + d(v_2)$ .
- (c) Every basis for a particular space have equal number of vectors and this number is called the dimension of the space.

**Solution:**

13. For the following set of statements, state whether they are True or False. Provide a valid reasoning or example/counter-example to justify your judgement.

- (a)  $A$  and  $A^T$  have the same left nullspace.
- (b)  $A$  and  $A^T$  have the same number of pivots.
- (c) If the row space equals the column space then  $A^T = A$ .
- (d) If  $A^T = -A$  then the row space of  $A$  equals the column space.

**Solution:**

14. Consider a plane  $x - 2y + 3z = 0$  in  $R^3$ . Find the basis vectors which span this plane. Further, find a basis for the following :-

- (a) intersection of this plane with the standard XY plane.
- (b) all vectors perpendicular to the plane.

**Solution:**

15. Which of the following vectors are linearly independent:

- (a) (1,0) and (1,0.001)
- (b) (2,1) and (0,0)
- (c) (-1,-1) and (1,1)
- (d) (5,0) and (0,5)

**Solution:**

16. Compute the rank and nullity of the given matrix  $M = \begin{bmatrix} 2 & 0 & -1 \\ 4 & 0 & -2 \\ 0 & 0 & 0 \end{bmatrix}$

**Solution:**