

# CS5691: Pattern recognition and machine learning

## Quiz - 3

Course Instructor : Prashanth L. A.

Date : Mar-22, 2019 Duration : 40 minutes

Name of the student :

Roll No :

**INSTRUCTIONS:** For short answer questions, you do not have to justify the answer. For the rest, provide proper justification for the answers. Please use rough sheets for any calculations *if necessary*. Please **DO NOT** submit the rough sheets. **DO NOT** use pencil for writing the answers.

### I. Short answer questions

1. Consider a two-class classification problem in a three-dimensional space, where the classes are linearly separable. Suppose the optimal separating hyperplane, i.e., one with highest margin, is given by

$$2x_1 + x_2 + x_3 - 3 = 0.$$

For each of the following training examples, identify the class label, and whether it is a support vector or not: ( $\frac{1}{2}$  mark for each table entry)

example	class label	support vector or not
(1, 1, 1)		
( $\frac{1}{2}$ , 1, 0)		
(1, 1, 2)		

2. Consider a classification dataset with points  $x_1 = -1, x_2 = 1, x_3 = 100$ , and corresponding class labels  $y_1 = -1, y_2 = +1, y_3 = +1$ . Let  $w_1 = 0$ , and  $w_2 = 1$ . Let  $L_1(w)$  and  $L_2(w)$  denote the square-loss, and log-loss, respectively, for any  $w \in \mathbb{R}$ . Recall that square-loss is employed in linear regression, and log-loss (or cross-entropy loss) in logistic regression. Which of the following statements is true? (1 mark)

- (a)  $L_1(w_1) \leq L_1(w_2), L_2(w_1) \leq L_2(w_2)$ .
- (b)  $L_1(w_1) \leq L_1(w_2), L_2(w_1) \geq L_2(w_2)$ .
- (c)  $L_1(w_1) \geq L_1(w_2), L_2(w_1) \leq L_2(w_2)$ .
- (d)  $L_1(w_1) \geq L_1(w_2), L_2(w_1) \geq L_2(w_2)$ .

Answer:

3. Consider the optimization problem for finding the maximum margin separating hyperplane, assuming that the classes are linearly separable:

$$\min_{w \in \mathbb{R}^d} W^T W \quad \text{subject to } y_i (W^T X_i + b) \geq 1, i = 1, \dots, n, \quad (1)$$

where  $\{(X_i, y_i), i = 1, \dots, n\}$  is the training dataset. Consider the following variant of the problem in (1):

$$\min_{W \in \mathbb{R}^d} W^T W \quad \text{subject to } y_i (W^T X_i + b) \geq 100, i = 1, \dots, n. \quad (2)$$

State whether the optimal hyperplane found by solving (1) is the same as that obtained by solving (2). (1 mark)

*Answer:*

4. Suppose we collect data about the number of hours spent per week by students of CS5691, and whether they passed or not. Let  $\{(x_i, y_i), i = 1, \dots, 86\}$  denote the dataset, where  $x_i$  is the hours spent by student  $i$ , and  $y_i$  is a boolean indicating whether the student passed or not. We perform logistic regression on this dataset, and obtain the following output: weight  $w = 1.5$ , and  $b = -4$ .

Using the logistic regression model, what is the probability that a student who studies 2 hours per week, passes the quiz? (1 mark)

*Answer:*

## II. A problem that requires a detailed solution

1. Consider a two-class one-dimensional classification dataset with points  $x_1 = -2, x_2 = 2, x_3 = 3$  having class label  $-1$ , and  $x_4 = 1$  with class label  $+1$ .

Answer the following:

- (a) Is the data linearly separable? (1/2 mark)
- (b) Consider the transformation  $\phi(x) = (x, x^2)$ . Form a two-dimensional dataset with inputs transformed using  $\phi$ .
- (a) Show that the transformed problem is linearly separable. (1/2 mark)
- (b) Find the maximum margin separating hyperplane in this transformed problem, either by solving the SVM optimization problem using KKT conditions, or a geometric argument. (1 1/2 marks)
- (c) What is the margin of the optimal hyperplane obtained above? (1/2 mark)
- (c) In the part above, suppose the transformation is  $\hat{\phi}(x) = (2x, 2x^2)$ . Compare the margin resulting from  $\hat{\phi}$  to that obtained using  $\phi$ . (1 mark)